DNA-TR-81-35

LATE DUST PEDESTAL RISE MECHANISMS

D. H. Sowle

C)

R. H. Christian

Mission Research Corporation

P.O. Drawer 719

Santa Barbara, California 93102

31 August 1981

Technical Report


CONTRACT No. DNA 001-81-C-0162

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

THIS WORK SPONSORED BY THE DEFENSE NUCLEAR AGENCY UNDER RDT&E RMSS CODE B345081466 J11AAXAX00007 H2590D.

E FILE COPY

Prepared for
Director
DEFENSE NUCLEAR AGENCY
Washington, D. C. 20305



0 1 12 13 000

BLANK PAGES IN THIS DOCUMENT WERE NOT FILMED

Destroy this report when it is no longer needed. Do not return to sender.

PLEASE NOTIFY THE DEFENSE NUCLEAR AGENCY, ATTN: STTI, WASHINGTON, D.C. 20305, IF YOUR ADDRESS IS INCORRECT, IF YOU WISH TO BE DELETED FROM THE DISTRIBUTION LIST, OR IF THE ADDRESSEE IS NO LONGER EMPLOYED BY YOUR ORGANIZATION.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM					
1 nue =	ESSION NO. 3. RECIPIENT'S CATALOG NUMBER					
	13 112					
4 TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED					
LATE DUST PEDESTAL RISE MECHANISMS	Technical Report					
	6. PERFORMING ORG. REPORT NUMBER					
7 AUTHOR(s)	MRC-R-657 B CONTRACT OR GRANT NUMBER(s)					
D. H. Sowle R. H. Christian	DNA 001-81-C-0162					
9 PERFORMING ORGANIZATION NAME AND ADDRESS Mission Research Corporation	10 PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS					
P.O. Drawer 719	Subtask JllAAXAX000-07					
Santa Barbara, California 93102	Subsusk of framewood of					
11 CONTROLLING OFFICE NAME AND ADDRESS Director	12. REPORT DATE					
Defense Nuclear Agency	31 August 1981					
Washington, D.C. 20305	34					
14 MONITORING AGENCY NAME & ADDRESS(If different from Controll	ing Office) 15. SECURITY CLASS (of this report)					
	UNCLASSIFIED					
1	15a. DECLASSIFICATION DOWNGRADING					
	SCHEDULE N/A					
16. DISTRIBUTION STATEMENT (of this Report)						
Approved for public release; distribution	unlimited.					
17 DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if	different from Report)					
18 SUPPLEMENTARY NOTES						
This work sponsored by the Defense Nuclear RDT&E RMSS Code B345081466 J11AAXAX00007 F	Agency under					
RUIAE RMSS COde B345081466 JITAAXAXUUUU/ F	123900.					
19 KEY WORDS (Continue on reverse side if necessary and identify by b	lock number)					
	iota number)					
Nuclear Lofted Dust Clouds						
Dust Pedestal						
20 ABSTRACT (Continue on reverse side if necessary and identify by	riber)					
The mechanism of buoyant rise is found to	be quantitatively consistent with					
readily available photographic data on the	late dust pedestal of events					
Wasp Prime and Climax. The term "late" re						
initial pedestal rise, and can be convenie model. One implication of buoyancy as a	driving force for late nedectal					
I rise is that the average bulk density in π	most or all of the dust pedestal					
is small compared to air density, that is,	small compared to 10^{-3} g/cm ³ .					

DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)
20. ABSTRACT (continued)
A number of mechanisms other than buoyancy were investigated as causes
for the late rise with discouraging results; these mechanisms are listed along with a brief description of the reasons each is untenable.
1

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

PREFACE

Major Leon A. Wittwer, USAF, brought the possibility of buoyancy as a driving force to our attention, based upon the resemblence between a late dust pedestal and the top of a layer of summer cumulus clouds.

resultan/
continued
contin

1

TABLE OF CONTENTS

Section	<u>on</u>	<u>Page</u>
PRI	EFACE	1
LIS	ST OF TABLES	3
1	INTRODUCTION AND SUMMARY	5
2	TERMINOLOGY	8
3	DEMONSTRATION OF EXCESS VERTICAL PEDESTAL VELOCITY	10
	Quiescent Flow	10
	Excess Velocity	11
	Wasp Prime Data	11
	Results of Wasp Prime Comparison	14
	Climax Data	15
4	COMPARISON OF EXCESS RISE VELOCITY TO THAT EXPECTED FROM BUOYANCY	17
	Expected Rise Rate of Outstanding Lumps	17
	Estimate of $\Delta \rho / \rho_a$	19
	Comparison of Observations to Buoyant Expectations	21
5	PHENOMENA WHICH CANNOT LIFT DUST ADEQUATELY	23
	Popcorning	23
	Steaming	24
	Reverse Percolation	24
	Suction	25
	Turbulence	25
RFI	FERENCES	27

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Measured feature trajectories for Wasp Prime.	12
2	Characteristic radii of Wasp Prime features.	14
3	Wasp Prime excess velocities.	15
4	Climax data.	16
5	Buoyant predictions compared to data.	22

SECTION 1 INTRODUCTION AND SUMMARY

A rapid survey of available literature and data was undertaken in an attempt to become familiar with the state of understanding of the dust pedestal, and to determine if any areas of phenomenology appeared ripe for pursuit, particularly matters which might contribute to electromagnetic propagation interference understanding. We quickly became engrossed in the mysteries of the nearly rectangular vertical profile of the pedestal, its sharp edge, long continuing rise, near universal occurrence over a wide range of conditions, and other phenomena more minor in impact and less universal in occurrence, but equally mystifying, such as the marvelous central sand column of some events. This report documents results of our efforts to understand only one of the above aspects of the pedestal, the lofting mechanism for its late slow rise. As an accidental byproduct of this investigation, the rise rate of the sand column, but not its existence, is explained.

Section 2 explains the terminology of those aspects of the dust pedestal which concern us.

In Section 3 it is shown that at late times a rather steady excess velocity, nearly vertical, is present in the flow field in addition to that anticipated due to the rising fireball or its shocks and rarefactions; at least this is so in the two cases investigated to date. These are Wasp Prime (3.2 KT at 225 m) and Climax (61 KT at 407 m). These shots were chosen for the initial study for two reasons: (1) burst height was low enough to raise a dust pedestal but high enough not to create a crater which would confuse data with ejecta, and (2) data on these two shots were readily available, in excellent pictures with accompanying dimensional information. 2 a,b

In Section 4 it is shown that excess rise velocities measured for identifiable features at the top of the dust pedestal are reasonably consistent with theoretical estimates based on buoyancy estimates and measured feature size.

From consistency of buoyancy as a driving mechanism follows the conclusion that at least at late times in the buoyant portion of the pedestal, bulk dust density must be too low to seriously interfere with the force balance. These forces are estimated to be caused by 5 percent to 50 percent air density depletion due to thermal expansion. Accordingly, dust bulk density must be less than 5 percent to 50 percent of air density.

Examination of the photographs show the features on which this low dust density conclusion is based to be comparable to the pedestal thickness in dimension. This implies local "buoyant thermal" or vortex flow fields which extend throughout the vertical thickness of the pedestal. Thus is implied, although not with the strength of the dust density limit above, that this limit on bulk density extends throughout most, if not all of the pedestal.


Calculations of Gutsche, ³ Lauer, ⁴ and Thompson ⁵ show that at such low levels of dust density the radar propagation problem is reduced to one of angular scintillation; signal blackout cannot be overriding. Scaling to probable threat level yield is problematical at the moment, although the required extrapolation in yield is no greater than that between the two events here investigated—and bulk density has the same limit for these cases.

Finally, Section 5 outlines considerations which lead to rejection of a short list of candidate phenomena as causes of the late dust rise. Most or all of these phenomena may, in fact, occur but none seems adequate singly or in concert to do more than initiate a low dust cloud. Some other

mechanism must raise it rapidly during the early phase in preparation for the buoyant late phase.

SECTION 2 TERMINOLOGY

A qualitative outline of the early, formative stage of the pedestal is illustrated on the left of the sketch. The dust cloud rises immediately behind the shock, or the precursor, if present. The rise is rapid, typically at velocities of tens to hundreds of meters per second until height H_S is reached within a second of shock passage. After this precise time the rise rapidly slows, by about



an order of magnitude, but continues for a long time: minutes. The slowing in rise velocity is so abrupt that Liner, et al. have modeled the rise velocity as discontinuous at H_s . Very roughly,

$$H_{s} \sim 100 [Y(MT)]^{1/3}$$
 meters . (1)

Prior to reaching H_s , dust laden material rises at a rate of order

$$v_{-} \sim \text{hundreds of meters/sec}$$
, (2)

after reaching H_s , material slows to

$$v_{+} \sim \text{tens of meters/sec}$$
 (3)

and continues to rise for several minutes; presumably it gradually slows to a stop. None of the above numbers should be taken as accurate; see Reference 1 for one set of self consistent parameter values.

An approximation to the above behavior occurs without radial dependence other than timing out to the maximum pedestal radius, R_p , which is typically at a distance of two to three fireball radii from ground zero. Time starts counting when the shock or its precursor reaches a given point. Actually, many believe that H_s increases with distance from ground zero out to R_p , where the process of pedestal creation abruptly ceases.


The late pedestal, shown on the right of the sketch, is characterized by a very uneven top surface composed of more or less random lumps with apparent outer scale size about equal to pedestal height. Many lumps in this structure can be identified in successive frames of motion pictures; thus their development and motion can be followed for analysis.

An exception to the random structure is the so called sand column. This feature is observed for burst heights greater than a fireball radius (of course it would be invisible at lower heights) and less than two or three fireball radii. The sand column is aptly named; it appears to be a column of dirt which forms at ground zero promptly after the shock strikes. Its radius is about half that of the fireball and remains roughly constant in time. The sand column shoots upward as a cylinder, eventually overtaking the fireball and taking on the appearance of a classic mushroom cloud stem, albeit a bit narrower stem than usual. We draw attention to it here because, despite its extraordinary appearance, its motion appears to be due to the same forces which control the remainder of the late dust pedestal.

SECTION 3 DEMONSTRATION OF EXCESS VERTICAL PEDESTAL VELOCITY

QUIESCENT FLOW

Under ideal conditions, after the shock has gone and its reverberations calmed, the flow field exterior to the fireball should be predictable from the rising motion of the fireball. The dominant motion at position, r, should be the sum of that due to a dipole field exterior to a rising sphere of radius, a, adequate to encompass the fireball or perhaps a bit larger and, because the ground surface is



impenetrable, an image dipole below the ground. These spheres and their relation to an observation point at (x,z) are shown in the sketch.

If the fireball, represented by a sphere of radius, a, rises through an element of height, dh, then the material at (x,z) should be displaced by

$$dx = \frac{3a^{3}x}{2} [(h-z)/r^{5} + (h+z)/r^{5}] dh$$
 (4)

$$dz = a^{3} / [(h-z)^{2} - x^{2}/2]/r^{5} - [(h+z)^{2} - x^{2}/2]/r^{15}/dh$$
 (5)

where
$$r^2 = (h-z)^2 + x^2$$
 and $(r^*)^2 = (h+z)^2 + x^2$. (6)

If one wishes to analyze the motion of a feature visible on a photograph, it is not certain that the feature is in the plane which passes through the burst point perpendicular to the observer. In that case, one must assume a value of azimuth angle, ϕ and set

$$x = x_a/\cos\phi \tag{7}$$

where x_a is the "apparent" or projected value of x. The equation of motion for the apparent x is obtained by multiplying the right hand side of (4) by $\cos \phi$.

Attempts to match the observed motion of lumps in the Wasp Prime and Climax late dust pedestals to these simple flows were unsuccessful, the observed vertical motion being significantly greater than expected.

EXCESS VELOCITY

To investigate the effect of a possible excess vertical velocity, a term is added to Equation 5 for $\,\mathrm{d}z$.

$$dz_e = u dt = u(dt/dh)dh = u/U dh$$
 (8)

where u is the excess velocity and U is fireball rise velocity.

WASP PRIME DATA

Wasp Prime was a 3.2 KT air drop exploded at 225 meters above the Nevada Proving Grounds as event 9 of Operation Teapot.

Features identifiable in successive plates of Dudziak's report 23 were measured by hand directly from the report. The (x,z) trajectories measured for these features are shown in Table 1 and Figure 1, which also shows the fireball center location and fireball radius. Feature number 3

Table 1. Measured feature trajectories for Wasp Prime. Trajectories measured from west station photos in Ref. 2a are shown for five features. Feature 3 is the sand column, feature 2 is known to have $\phi \! = \! 0$. The first column has time in seconds post shot. Coordinates x and z are apparent Cartesian coordinates in meters, projected into the photographic plane. Note that features 1 and 2 are at negative values of x.

	Feature 1		Feature 2		Feature 3		Feature 4		Feature 5	
t(sec)	-x	Z	-x	Z	х	Z	x	Z	х	Z
1.005									222.3	12
1.095									222.8	11
1.298									218.8	13
1.503	124.2	19					130.8	15	210.8	13
1.698	122.3	20					133.7	18	207.7	17
1.903	120.3	22	80.3	18			131.7	20	203.7	19
2.001	120.8	22	76.8	19	0	55	131.2	20	205.2	19
2.205	119.4	23	74.4	22			127.6	20	205.6	20
2.400	118.1	26	75.1	25			127.9	23	206.9	22
2.498	116.2	28	74.2	26			127.8	25	204.8	24
2.700	114.1	28	72.2	27			127.8	26	206.8	24
2.904	115.4	29	72.4	27			123.6	27	202.6	25
3.002	113.0	30	69.0	28	0	95	125.0	28	204.0	26
3.099	112.0	31	71.0	30			125.0	30	204.0	25
3.505	110.1	32	71.1	32			122.9	28	200.9	25
3.807	108.7	33	69.7	35			120.3	29	200.3	26
4.008	104.8	37	64.8	39	0	132	124.2	30	203.2	27
4.25	100.7	35	65.7	40			121.3	30	201.3	28
4.50	100.7	37	67.7	42			119.3	31	199.3	28
4.75	97.8	40	65.8	45			119.2	33	202.2	28
5.00	99.8	40	67.8	47	0	175	116.2	33	197.2	30
5.50							117.1	33	195.1	30
6.00							118.7	35	198.7	34
6.50						;	120.6	37.5	195.6	33




Figure 1. Motion of Wasp Prime features, as measured from Ref. 2a. Encircled integers are times in seconds when the features, identified numerically in diamonds below the horizontal axis, were located at circle centers. The quantity af, is apparent horizontal fireball radius.

is the top of the sand column. Feature number 2 is identifiable from the north station as well as the west station and, as viewed from west station, has $\phi = 0$. The apparently erratic motion of the features is not believed to be real, but rather is due to crude measuring techniques; nevertheless, the trend of motion should be reliable.

Later we will have need of the characteristic radius of each feature. Approximate values measured during the observation interval are shown in Table 2.

Table 2. Characteristic radii of Wasp Prime features.

RESULTS OF WASP PRIME COMPARISON

Equations 4 through 8 were entered into a computer, along with an interpolated tabular fit to fireball center position, h, and radius, $a_{\mathbf{f}}$. A number of ratios of equivalent sphere radius, a, to horizontal fireball radius, $a_{\mathbf{f}}$, were attempted. The quality of obtainable fits to the five observed features deteriorated markedly for $a/a_{\mathbf{f}}$ outside the range

$$1.2 \leq a/a_{f} \leq 1.3$$
 (9)

This range is determined basically by the change in $\, x \,$ over the observation time. Smaller values of $\, a/a_{\hat{f}} \,$ gave estimates of motion too small and larger values gave estimates too large. Our first conclusion, then, is that the effective value of radius for the volume of air moving with the fireball is 1.2 to 1.3 times the observed horizontal fireball radius.

The second conclusion reached is that significant upward velocites in excess of that due to the ideal flow field are necessary for each feature. The best time averaged value of excess velocity, u, is listed for each feature in Table 3, taking the mean fireball rise velocity during the period of observation to be 40 m/sec.

Best fits for features 4 and 5 occur at values of \cos ; of 0.7 to 0.8, while $\cos \phi = 1.0$ worked best for the others.

Table 3. Wasp Prime excess velocities

Feature		1	2	3	4	5
u	a/a _f =1.2	3.2	4.8	22.	4.0	4.8
(m/s)	a/a _f =1.3	4.0	5.6	18.	4.4	4.8

CLIMAX DATA

Climax was a 61 kt air drop exploded 407 meters above the Nevada Proving Ground as event 11 of series Upshot-Knothole. It was one of the largest yield devices ever tested over the Nevada Proving Ground. The scaled height of burst $(\Upsilon^{1/3})$ is about two-thirds that of event Wasp Prime.

Experience with Wasp Prime showed that most of the vertical motion was in fact considerably in excess of that due to an ideal flow field. Thus elaborate computations of dipole and image dipole flow fields were not really necessary to extract an approximate value of excess vertical velocity. Only a fairly small correction for the effect of the dipole fields is necessary. Accordingly, values of observed upward velocities for event Climax were corrected for ideal flow field effects by factors found in analysis of Wasp Prime data. For all points other than the sand column, 80 percent of the observed velocity was taken to be excess. For the sand column, 45 percent of observed velocity was taken to be excess.

A summary of Climax features that were analyzed based on photographic data and scales provided in Reference 2b appears in Table 4.

Table 4. Climax data. The first column is feature identification, the second column approximate feature horizontal location, x(m) at about 8 seconds post shot as viewed from the west station. The third and fourth columns show the observed total velocity, $V_Z(m/s)$, and the estimated excess upoward velocity, u(m/s). The fifth column is the feature's characteristic radius in meters. Feature 3 is the sand column.

Feature	x(m)	V _z (m/s)	u(m/s)	R(m)
1	-600	10.7	8.6	36
2	-350	15.9	12.7	55
3	0	80.0	36.0	128
4	800	10.0	8.0	30.3

SECTION 4 COMPARISON OF EXCESS RISE VELOCITY DATA TO THAT EXPECTED FROM BUOYANCY

Consideration of the general appearance of a late pedestal and of the amount of energy striking the area as thermal radiation or shock leads one to suspect that the dust pedestal may be buoyant due to thermal expansion of heated air, unless in fact the air is so laden with dust as to overpower this buoyancy. In order to investigate this possibility, a comparison of the excess rise velocity observed for individual features can be made to that expected for buoyantly driven rise.

As a preliminary observation, note that the density of dust necessary to record photographically is relatively small and comes almost entirely from the finest dust particles in the distribution. For example, dust density of the order of 10^{-5} to 10^{-6} gm/cm³ consisting of 10 micron particles, too thin to interfere seriously with many radio or radar frequencies, would be sufficient to record on film as an opaque cloud.

EXPECTED RISE RATE OF OUTSTANDING LUMPS

If the pedestal material rises because dust in combination with warmed, expanded air is buoyant then the objects measured, which must strongly protrude to be observable, should behave roughly as rising hot air "thermals" or buoyant spheres.

An estimate of the buoyant rise velocity may be obtained by equating the buoyant force to the drag force and solving for the quasi-steady velocity. The buoyant force, by Archimedes principle, is the difference between the weight of the displaced fluid and the weight of the body. For

this purpose we assume a spherical bubble of radius R,

$$F_{b} = \frac{4\rho}{3} R^{3} (r_{a} - \rho) g \tag{10}$$

where ρ_a is the density of ambient air, ρ is the (average) density of the bubble, and g is gravitational acceleration.

The drag force of an object of cross sectional area, πR^2 , moving at a velcoity, u, is by definition,

$$F_{d} = C_{D} \rho_{a} \frac{u^{2}}{2} (\pi R^{2})$$
 (11)

where the drag coefficient is a function of body shape and Reynolds number. We will use a value of $C_{\rm D}$ = 0.6, appropriate to a smooth sphere in high Reynolds number, laminar flow.* Equating these forces and solving for u yields

$$u = \left[\frac{8}{3C_D} g R \frac{\Delta \rho}{\rho_a} \right]^{1/2} \approx 2.1 \sqrt{g R \Delta \rho/\rho_a} , \qquad (12)$$

where $\Delta \rho = \rho_a - \rho$. This quasi-equilibrium estimate neglects impulsive forces necessary during the initiation of the motion and therefore is an upper limit.

The smallest likely rise rate may be obtained by extrapolating a formula due to McCartor and Wortman. For the early rise rate of a perfect, spherical, hollow bubble in quiescent air they find

$$u \approx 0.8 \sqrt{gR} \qquad . \tag{13a}$$

To account for the actual, but relatively low, density of the pedestal features we modify their formula to include the relative density, i.e.,

$$u = 0.8 \sqrt{gR\Delta\rho/\rho_a} . (13b)$$

The decrease in $C_{\rm D}$ that occurs for a solid sphere at Reynolds number greater than 10^4 is due to detachment. For underdense gas spheres, this seems unlikely to us.

From the above two limits we conclude that buoyant forces should lead to a feature rise velocity in the range

$$0.8 \lesssim \frac{u}{\sqrt{gR\Delta\rho/\rho_a}} \lesssim 2.1 \qquad . \tag{14}$$

ESTIMATE OF Δρ/ρa

Ganong and Whitaker 8 use several empirical but reasonable formulae in an attempt to match air temperature measurements near the surface following explosions of the Tumbler series, and also to match the precursor shock behavior of shot Priscilla. Based on comparison of these several formulae to Jata, an appropriate formula for bomb thermal radiation which is deposited in air as thermal energy, ϵ , appears to be

$$\varepsilon = \frac{f_c f_r W h}{4\pi r^3}$$
 (15)

where f_c is bomb thermal to air thermal coupling fraction, f_r is bomb thermal radiation fraction, W is total yield, h is burst height above ground zero, and r is the distance from burst point to surface, approximately

$$r = (h^2 + x^2)^{1/2} (16)$$

where as before, x is horizontal distance from ground zero. Ganong and Whitaker found best agreement with precursor data in our range of x for $f_c = 1/6$.

Equation 15 provides a means of estimating the energy deposited per unit area. It is then necessary to discover, or guess, how this energy mixes vertically into a volume which must eventually expand to pressure equilibrium with the ambient atmosphere. All evidence points out to very rapid initial mixing as evidenced by the fast initial rise of the pedestal. Leaning heavily on this sparce evidence, we assume the energy deposited

in some area to be uniformly mixed vertically to a height, ℓ_0 , so quickly that no net work is done on the exterior during this expansion via mixing, thus no energy is lost.

The resultant mix of air at ambient density and uniform superambient temperature is then assumed to expand adiabatically to pressure equilibrium.

When energy ϵ per unit area is distributed in air to height, ℓ_o , at ambient density, ρ_a , the air specific internal energy will be

$$I_o = \frac{\varepsilon}{\ell_o \rho_a} + I_a \tag{17}$$

where I_a is ambient specific internal energy,

$$I_a = \frac{p_a}{(\gamma - 1)\rho_a} \tag{18}$$

where γ is the normal air polytropic constant.

Then, substituting (18) into (17), the raised pressure, p_0 , will be given by

$$\frac{\mathbf{p_o}}{\mathbf{p_a}} = \frac{\mathbf{I_o}}{\mathbf{I_a}} = 1 + \frac{(\gamma - 1)\varepsilon}{\ell_o \mathbf{p_a}} \tag{19}$$

and, after adiabatic expansion back to ambient pressure, $\,p_a^{}$, the density will be given by

$$\frac{\rho}{\rho_a} = \left(\frac{P_o}{P_a}\right)^{1/\gamma} = \left[1 + \frac{(\gamma - 1)\varepsilon}{\ell_o P_a}\right]^{1/\gamma} \tag{20}$$

but, since expansion is almost entirely one dimensional in the vertical direction, raising material at height ℓ_0 to the pedestal height, H_s , at the end of the rapid rise, it is also true that

$$\frac{\rho}{\rho_{\mathbf{a}}} = \frac{l_{\mathbf{o}}}{H_{\mathbf{S}}} \qquad . \tag{21}$$

Solving (21) for ℓ_{o} and substituting ℓ_{o} into (20) yields

$$\frac{\rho}{\rho_a} = \left[1 + \frac{(\gamma - 1)\varepsilon}{H_s p_a} \frac{\rho_a}{\rho}\right]^{-1/\gamma} . \tag{22}$$

The transcendental Equation (22) can be solved for $\left. \rho/\rho_{a}\right.$ and hence the desired quantity,

$$\Delta \rho / \rho_{\mathbf{a}} = 1 - \rho / \rho_{\mathbf{a}} \quad . \tag{23}$$

COMPARISON OF OBSERVATIONS TO BUOYANT EXPECTATIONS

Values of parameters for use in evaluation of (14), (15), (16), and (23) are

$$g = 9.8 \text{ m/sec}^2$$
, $f_r = 0.4$, $\gamma = 1.4$, and $p_a = .8 \times 10^6 \text{ dynes/cm}^2$.

For Wasp Prime, Y = 3.2 kt , h = 225 m, $H_s \approx 20$ m .

For Climax, Y = 61 kt , h = 407 m, $H_S \approx 50$ m .

Evaluations of pertinent quantities and comparisons to measurements are shown in Table 5. The coupling fraction, $f_{\rm c}$, was varied from slightly less than the value Ganong and Whitaker chose to nearly twice their value.

Table 5. Buoyant predictions compared to data. The value of u shown for Wasp Prime is the average of those shown in Table 3 for the two values of a/a_f .

Event	Feature	x(m)	R(m)	u(m/s)	ε(erg/cm ²)	∆ \0	u /∕Rg∆o/p	Δρ/ρ_	u /√Rg Δρ / ρ
Wasp	1	-124	5.9	3.6	1.80×10^{9}	.31	.85	.16	1.2
	2	-80	4.4	5.2	2.36×10^9	.38	1.3	.20	1.8
Prime	3	0	44	20	2.82 × 10 ⁹	.44	1.5	.24	2.0
1 7 mic	4	131	5.9	4.2	1.82×10^9	.30	1.0	.16	1.4
	5	222	7.3	4.8	1.02 × 10 ⁹	.18	1.3	.09	1.9
	1	-660	36	8.6	2.90 × 10 ⁹	.20	1.0	.10	1.5
C1imax	2	-350	55	12.7	7.15×10^9	.45	0.82	.24	1.1
OT THIOX	3	0	128	36.0	1.64×10^{10}	.79	1.0	.50	1.4
	4	800	30	8.0	1.53×10^9	.11	1.4	.05	2.1
$f_c = 0.30$							0	f _c	= 0.15

Inspection of columns 8 and 10 in Table 5 shows that observed values of excess velocity, taken from Tables 3 and 4, fall within the range predicted by Equation 14. Thus, qualitatively and quantitatively, the data are consistent with buoyancy as the major driving force at late times, to the accuracy of measurements and within the confidence possible with an incomplete theory.

SECTION 5 PHENOMENA WHICH CANNOT LIFT DUST ADEQUATELY

This section presents a partial list of phenomena which might be suggested as candidates for raising dust to the early pedestal height, $\rm H_S$, or for causing the late time pedestal rise. None of them singly can explain observations; in fact, it seems most unlikely that all of them in concert can explain either early pedestal rapid rise to $\rm H_S$ or late slow rise. This is not to say that these phenomena are absent nor, as a member in some possible chain of events, unimportant. For example, popcorning is certainly real and may provide the dust, at a very low level, which a Hess bubble or some other mechanism lifts to $\rm H_S$, but popcorning itself does not raise dust as high as $\rm H_S$.

POPCORNING

This mechanism has been popular at least since the early 1950's. Basically, the mechanism consists of flash heating water laden naterial via bomb thermal, neutrons or gamms so severely that steam forms explosively causing dust to fly up into the air.

From (15) the maximum energy available at, say, x = h, is

$$_{M} = \frac{f_{c}f_{r}W}{\sqrt{2}8\pi\hbar^{2}} \tag{23}$$

from bomb thermal; much less is available from gammas or neutrons.

Taking the energy necessary to raise water to the boiling point and convert it to steam as I_s , ergs/gm, the amount of steam possible is

$$\mathfrak{m} = \frac{\mathbf{f_c f_r W}}{\sqrt{2} 8\pi \mathbf{I_s h^2}} \qquad . \tag{24}$$

Using Climax as an example and taking f_r = 0.4, f_c = 1/3, I_s = 2.5 × 10 10 erg/gm.

$$m = 0.25 \text{ gram/cm}^2$$
.

If this steam then expands to density comparable to ambient air, it will reach up to height

$$H_p = m/10^{-3} = 250 \text{ cm} = 2.5 \text{ m}$$
 (25)

about a factor of 20 short of the required rise.

STEAMING

Here one imagines the shock, perhaps with help from bomb thermal radiation, heating the ground and creating steam, or some other outgased product, at a rate slow compared to popcorning. Dust is presumed carried upward with the gas. Steaming fails on the same basis that procorning fails; even less energy is available in the shock than in the thermal pulse.

REVERSE PERCOLATION

In this process air is forced into soil cavities by the high pressure forward portion of the shock, then air plus soil is blasted upward when the pressure gradient reverses on the back side of the positive phase. Zernow and coworkers¹⁰ have shown the process to be effective in lifting dessicated sand tens of centimeters under fairly realistic conditions. By "fairly" is meant a basically realistic pressure profile and hopefully realistic lower boundary condition (both are critical) but overoptimistic in that the material is: (1) dessicated, and (2) uncompacted.

Based upon their experiments, the process does not seem capable of raising dust more than a few meters at most. Further, their experiment showed the process to fail completely in very wet sand, whereas both Teapot 12 (Met) and Tumblers 1 through 4 show no effect on the dust pedestal of either artificial or natural wetting down of the shot region relative to dry desert surface.

SUCTION

Suction is due to the rarefaction behind the shock and can be viewed as a continuation of reverse percolation. After the positive phase of the shock has passed, a rarefaction, typically a 20 percent reduction in atmospheric pressure, exists over a region similar in extent to the pedestal. If air at ambient pressure exists in soil interstitial regions it may be expelled and may carry some dust with it. Any rational perturbation of linear to nonlinear D'Arcy law interstitial flow yields only about a meter of air lifted by this mechanism.

TURBULENCE

Turbulence has been suggested both as a mechanism for the initial pedestal rise and separately as a mechanism for the late rise.

As a mechanism to drive the early rapid rise turbulence seems not to withstand scrutiny on a rather basic level. In the most extreme case the shock proceeds outward and the pedestal proceeds upward at almost the same rate of speed. If the pedestal were rising as a consequence of some random process, which must be ultimately driven by the shock, it would rise with a \sqrt{t} law, or rise some random fraction of the time—but could not rise always and steadily at a rate near that of shock propagation.

furbulence as a driver of the late pedestal rise is not unlikely in principle, but it does not fit the observations. Of the nine pedestal features whose motions we followed (make it seven if one wishes to consider the sand columns to be special cases) all had excess velocities which were not only in the same general direction, a result most unlikely for a turbulent velocity field, but were quantitatively smooth and rather well predicted, a result nigh impossible in a random flow field.

REFERENCES

- 1. Liner, R. T., et al., <u>Nuclear Precursor Phenomenology and Sweep-up Dust Cloud Model</u>, DNA 3781F, Science Applications, Inc., McLean, Virginia, 4 November 1975 (U).
- 2a. Dudziak, W. F., et al., Photographic Atlas Event WASP PRIME, DNA 3576F, 9 May 1975 (U).
- 2b. Dudziak, W. F., et al., Photographic Atlas of Event CLIMAX, Unpublished.
- 3. Gutsche, S. L., X-Band Attenuation from a Nuclear Dust Cloud Pedestal:

 Boundary Calculations and Determination of Sensitivity to Key Parameters,

 MRC-N-464, 15 May 1981 (U).
- 4. Lauer, C. J., Angle of Arrival Fluctuations in a Nuclear Dust Cloud Pedestal, MRC-N-476, August 1981 (U).
- 5. Thompson, James H., <u>Preliminary Estimates of the Propagation Effects in a Nuclear Pedestal Region</u>, <u>Presentation at LoAD Dust Task Meeting</u>, <u>Defense Nuclear Agency</u>, <u>Washington</u>, D.C., 11 March 1981 (U).
- 6. See Lamb's <u>Hydrodynamics</u>, Sixth Edition, Dover Press, p. 161, 214 or <u>Physics</u> of <u>High Altitude Nuclear Burst Effects</u>, DNA 4501F, p 480, December 1977 (U).
- 7. McCartor, G. D., and W. R. Wortman, Unpublished calculation, Mission Research Corporation.
- 8. Ganong, G. P., and W. A. Whitaker, The Nuclear Blast Precursor, AFWL-TR-69-199, Vol. 1, March 1969 (U).
- 9. Hess, R. V., Interaction of Moving Shocks and Hot Layers, Langley Aeronautical Laboratory, NACA Technical Note 4002, May 1957 (U).
- 10. Zernow, L., et al., An Experimental Study of the 'Reverse Percolation' Lofting Process in a Sand Medium, DNA 3210F, Shock Hydrodynamics Division of Whittaker Corporation, 16 November 1973 (U).

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE DEPARTMENT OF THE ARMY (Continued) Assistant Secretary of Defense Comm, Cmd, Cont & Intell ATTN: Dir of Intelligence Sys, J. Babcock BMD Systems Command Department of the Army 2 cy ATTN: BMDSC-HW Command & Control Technical Ctr U.S. Army Communications R&D Command Department of Defense ATTN: DRDCO-COM-RY, W. Kesselman ATTN: C-312, R. Mason ATTN: C-650, G. Jones U.S. Army Comm-Elec Engrg Instal Agency ATTN: CCC-CED-CCO, W.Neuendorf ATTN: CCC-EMEO-PED, G. Lane 3 cy ATTN: C-650, W. Heidig Defense Intelligence Agency ATTN: Dir ATTN: DB-4C, E. O'Farrell ATTN: DB, A. Wise ATTN: DT-1B ATTN: DC-7B U.S. Army Foreign Science & Tech Ctr ATTN: DRXST-SD U.S. Army Materiel Dev & Readiness Cmd ATTN: DRCLDC, J. Bender Defense Nuclear Agency ATTN: NAFD ATTN: STNA U.S. Army Missile Intelligence Agency ATTN: YSE, J. Gamble ATTN: RAEE U.S. Army Nuclear & Chemical Agency ATTN: NATD ATTN: Lib 4 cy ATTN: TITL 6 cy ATTN: RAAE DEPARTMENT OF THE NAVY Office of Naval Research Defense Tech Info Ctr ATTN: Code 420 ATTN: Code 465 12 cy ATTN: DD ATTN: Code 421 Field Command Defense Nuclear Agency ATTN: FCPR, J. T. McDaniel DEPARTMENT OF THE AIR FORCE Air Force Geophysics Lab Field Command ATTN: OPR, H. Gardiner Defense Nuclear Agency ATTN: OPK, H. Gardiner ATTN: OPR-1 ATTN: LKB, K. Champion ATTN: OPR, A. Stair ATTN: S. Basu ATTN: PHP ATTN: PHI, J. Buchau ATTN: R. Thompson Livermore Branch ATTN: FCPRL Interservice Nuclear Weapons School ATTN: TTV Under Secretary of Defense for Rsch & Engrg Department of Defense Air Force Weapons Lab ATTN: Strategic & Space Sys (OS) Air Force Systems Command ATTN: SUL ATTN: NTYC ATTN: NTN National Security Agency ATTN: W-32, O. Bartlett ATTN: B-3, F. Leonard ATTN: R-52, J. Skillman Air University Lib Department of the Air Force DEPARTMENT OF THE ARMY ATTN: AUL-LSE Assistant Chief of Staff for Automation & Comm Department of the Army ATTN: DAAC-ZT, P. Kenny Space Division Department of the Air Force ATTN: SKA, D. Bolin ATTN: SKY, C. Kennedy Atmospheric Sciences Lab U.S. Army Electronics R&D Command ATTN: DELAS-EO, F. Niles Space Division Department of the Air Force 2"D Advanced Technology Ctr ATTN: YGJB, W. Mercer

Department of the Army ATTN: ATC-O, W. Davies ATTN: ATC-T, M. Capps

OTHER GOVERNMENT AGENCIES

Central Intelligence Agency ATTN: OSWR/NED

Department of Commerce National Bureau of Standards ATTN: Sec Ofc for R. Moore

Department of Commerce National Oceanic & Atmospheric Admin ATTN: R. Grubb

Institute for Telecommunications Sciences National Telecommunications & Info Admin

ATTN: W. Utlaut ATTN: A. Jean ATTN: L. Berry

DEPARTMENT OF ENERGY CONTRACTORS

Los Alamos National Lab ATTN: P. Keaton ATTN: R. Taschek ATTN: G-6, E. Jones ATTN: C. Westervelt ATTN: D. Simons ATTN: MS 664, J. Zinn ATTN: MS 670, J. Hopkins

DEPARTMENT OF DEFENSE CONTRACTOR

Aerospace Corp ATTN: T. Salmi ATTN: J. Straus ATTN: R. Slaughter ATTN: V. Josephson ATTN: I. Garfunkel ATTN: D. Olsen ATTN: N. Stockwell ATTN: S. Bower

Analytical Systems Engineering Corp ATTN: Radio Sciences

Analytical Systems Engineering Corp ATTN: Security

Barry Research Corp ATTN: J. McLaughlin

BDM Corp ATTN: L. Jacobs ATTN: T. Neighbors

Berkeley Research Associates, Inc ATTN: J. Workman

BETAC ATTN: J. Hirsch

Boeing Co

ATTN: M/S 42-33, J. Kennedy ATTN: G. Hall ATTN: S. Tashird

Booz-Allen & Hamilton, Inc ATTN: B. Wilkinson

Communications Satellite Corp ATTN: D. Fang

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

University of California at San Diego Electrical Engineering Computer Sciences ATTN: H. Booker

Charles Stark Draper Lab, Inc ATTN: J. Gilmore ATTN: D. Cox

Computer Sciences Corp ATTN: F. Eisenbarth

Comsat Labs ATTN: G. Hyde

Cornell University Department of Electrical Engineering ATTN: D. Farley, Jr ATTN: M. Kelly

E-Systems, Inc Center for Planning & Analysis ATTN: R. Berezdivin

Electrospace Systems, Inc ATTN: H. Logston

ESL, Inc ATTN: J. Marshall

General Electric Co ATTN: A. Harcar

General Electric Co ATTN: A. Steinmayer ATTN: C. Zierdt

General Electric Co ATTN: F. Reibert

General Electric Co ATTN: G. Millman

General Research Corp ATTN: J. Ise, Jr ATTN: J. Garbarino

Harris Corp ATTN: E. Knick

Horizons Technology, Inc ATTN: R. Kruger

HSS, Inc ATTN: D. Hansen

IBM Corp ATTN: F. Ricci

University of Illinois ATTN: Security Supr for K. Yeh

Institute for Defense Analyses ATTN: J. Aein
ATTN: H. Gates
ATTN: H. Wolfhard
ATTN: E. Bauer

JAYCOR

ATTN: J. Sperling

International Tel & Telegraph Corp Mitre Corp ATTN: W. Rice ATTN: Tech Lib ATTN: M. Horrocks ATTN: J. Wheeler ATTN: W. Hall **JAYCOR** ATTN: W. Foster ATTN: J. DonCarlos Pacific-Sierra Research Corp ATTN: E. Field, Jr ATTN: F. Thomas ATTN: H. Brode Johns Hopkins University ATTN: T. Potemra ATTN: J. Phillips ATTN: T. Evans ATTN: J. Newland ATTN: P. Komiske Pennsylvania State University ATTN: Ionospheric Research Lab Photometrics, Inc ATTN: I. Kofsky Kaman Tempo ATTN: T. Stephens ATTN: DASIAC Physical Dynamics, Inc ATTN: W. McNamara ATTN: E. Fremouw ATTN: W. Knapp Physical Research, Inc ATTN: R. Deliberis Linkabit Corp ATTN: I. Jacobs Litton Systems, Inc ATTN: R. Grasty R & D Associates ATTN: R. Lelevier ATTN: R. Turco Lockheed Missiles & Space Co, Inc ATTN: H. Ory ATTN: R. Johnson ATTN: W. Imhof ATTN: B. Gabbard ATTN: M. Gantsweg ATTN: W. Wright ATTN: C. Greifinger ATTN: W. Karzas ATTN: M. Walt Lockheed Missiles & Space Co, Inc ATTN: Dept 60-12 ATTN: D. Churchill ATTN. C. Old ATTN: F. Gilmore ATTN: P. Haas R & D Associates ATTN: B. Youn M.I.T. Lincoln Lab ATTN: D. Towle ATTN: E. Bedrozian Martin Marietta Corp ATTN: R. Heffner ATTN: C. Crain Riverside Research Institute ATTN: V. Trapani McDonnell Douglas Corp ATTN: J. Moule ATTN: G. Mroz ATTN: W. Olson ATTN: N. Harris Rockwell International Corp ATTN: R. Buckner ATTN: R. Halprin Rockwell International Corp Meteor Communications Consultants ATTN: S. Quilici ATTN: R. Leader Santa Fe Corp ATTN: D. Paolucci Mission Research Corp ATTN: S. Gutsche ATTN: R. Bogusch Science Applications, Inc ATTN: L. Linson ATTN: D. Hamlin ATTN: E. Straker ATTN: C. Smith ATTN: D. Sappenfield ATTN: F. Fajen ATTN: Tech Lib ATTN: R. Hendrick ATTN: R. Filb 4 cy ATTN: D. Sowle Science Applications, Inc 4 cy ATTN: R. Christian ATTN: SZ 5 cy ATTN: Doc Con Science Applications, Inc Mitre Corp ATTN: J. Cockeyne ATTN: A. Kymmel Technology International Corp ATTN: W. Boquist ATTN: C. Callahan ATTN: G. Harding ATTN: B. Adams

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

SRI International
ATTN: W. Chesnut
ATTN: R. Leadabrand
ATTN: D. Neilson
ATTN: J. Petrickes
ATTN: R. Livingston
ATTN: G. Price
ATTN: W. Jaye
ATTN: C. Rino
ATTN: A. Burns ATTN: A. Burns ATTN: R. Tsunoda

ATTN: G. Smith ATTN: M. Baron

Sylvania Systems Group ATTN: R. Steinhoff ATTN: I. Kohlberg ATTN: J. Concordia

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Tri-Com, Inc ATTN: D. Murray

TRW Defense & Space Sys Group ATTN: R. Plebuch ATTN: D. Dee

Utah State University

ATTN: K. Baker ATTN: J. Dupnik ATTN: L. Jensen

Visidyne, Inc ATTN: C. Humphreys ATTN: J. Carpenter